

# Effect of Alcohol Extracts of *Juniperus Phoenicea* and *Crataegus azarolus* on Diabetes Induced by Streptozotocin in Experimental Rats

Neveen M. Zeima<sup>1</sup>

1Nutrition and Food Science Dept., Faculty of Home Economics

Al-Azhar University, Egypt

[neveenzeima@azhar.edu.eg](mailto:neveenzeima@azhar.edu.eg)



## مجلة البحوث في مجالات التربية النوعية

معرف البحث الرقمي DOI: 10.21608/JEDU.2023.206055.1866

المجلد التاسع العدد 46 . مايو 2023

التقييم الدولي


P-ISSN: 1687-3424

E- ISSN: 2735-3346

موقع المجلة عبر بنك المعرفة المصري <https://jedu.journals.ekb.eg/>

موقع المجلة <http://jrfse.minia.edu.eg/Hom>

العنوان: كلية التربية النوعية . جامعة المنيا . جمهورية مصر العربية





## Effect of Alcohol Extracts of *Juniperus Phoenicea* and *Crataegus azarolus* on Diabetes Induced by Streptozotocin in Experimental Rats

### Abstract

Diabetes mellitus (DM) includes a diverse range of diseases associated with hypo- or insensitive insulin secretion. Thus, the current study aimed to examine the antihyperglycemic effects of alcoholic extract of leaves of *Juniperus Phoenicea* and *Crataegus azarolus* leaves in streptozotocin-stimulated diabetic rats (STZ, intraperitoneal, 65 mg kg<sup>-1</sup>). Thirty six rats (*Sprague-Dawley*) that weights (200±10 g) were divided randomly into six equal groups. Group (1) and group (2) were kept as negative and positive controls (- and +), respectively. Group (3) and group (4) were orally given the *Juniperus Phoenicea* extract in 200 and 400 mg kg<sup>-1</sup> B.wt .day<sup>-1</sup> doses, respectively, whereas group (5) and group (6) were orally given the *Crataegus azarolus* extract in 200 and 400 mg kg<sup>-1</sup> B.wt day<sup>-1</sup> doses, respectively, for a month. Finally, when the experiment was ended, a biological evaluation was determined. Blood glucose, serum insulin, heart enzymes, liver function, lipid profile, kidney function, and TNF- α were estimated in serum. In pancreas tissue, GSH, SOD and MDA were evaluated, as well as the chemical composition of herbs. The results exhibited that the alcoholic extract of *Juniperus Phoenicea* and *Crataegus azarolus* leaves significantly increased the biological evaluation, normalized serum levels of liver functions, kidney function, enhanced the lipid profile, lessened blood glucose, heart enzymes and MDA and increased levels of insulin, GSH and SOD than those of control diabetic rats. This work suggests that *Juniperus Phoenicea* and *Crataegus azarolus* leaves extracts administration can decline blood glucose levels and the occurrence of various problems resulting from hyperglycemia.

**Key words:** *Juniperus Phoenicea*; *Crataegus azarolus*; Streptozotocin; Blood glucose; Antioxidant enzymes.

## Introduction

Diabetes is a severe metabolic illness distinguished by insulin production, which leads to hyperglycemia and irregularities in protein metabolism, carbohydrates, and lipids (Nacer *et al.*, 2020). Diabetes is classified as a chronic irregular metabolic illness by the World Health Organization. Consequently, DM can be classed into two categories: type 1 and 2. Type 1 results from the immune system's dysfunctional performance, resulting in  $\beta$  cells destruction in islets and a disorder in insulin release. In contrast, type 2 is caused by insulin resistance and insufficient insulin secretion because of unhealthy lifestyle choices and thus obesity (Huang and Chen, 2023). Regardless of an individual's age, sex, socio-economical background, or race, diabetes incidence is steadily rising (Okoduwa *et al.*, 2017). Diabetes influences more than four hundred million people throughout the world, it is also supposed to increase this number to more than 600 million by 2040, with most of them having type 2 diabetes with huge disturbances in insulin production or its signal transmission (Hassan *et al.*, 2021). Also, it is associated with micro- and macro-vascular issues that adversely impact the functioning of several organs, such as retinopathy, cardiovascular disease, nephropathy, and others (Hacioglu *et al.*, 2021). Due to the higher oxidation rate of glucose in mitochondria, hyperglycemia has been shown to promote the producing of too many reactive O species (ROS) in people with diabetes (Al-Brakati *et al.*, 2020). Elevated levels of ROS and inhibition the defense system of antioxidants may cause increased lipid peroxidation, oxidative injury, insulin resistance, and cell injury (Hacioglu *et al.*, 2021). Also, diabetes might speed up the lipolysis process, which raises the level of free fatty acids, promoting the release of cytokines including interleukin-6 in addition to factor- $\alpha$  of tumor necrosis. The latter stimulates caspase transmission, that is crucial in inducing the death of cells through apoptosis (Giribabu *et al.*, 2017).

Diabetes has been the subject of much study, along with the creation of effective anti-diabetic medicines. Unfortunately, these medicines cannot restore the aforementioned diabetic problems, and their long-term use may result in medicine resistance and severe consequences such as acute renal damage (Chaudhary *et al.*, 2016 and Luo *et al.*, 2021). Many phyto-extracts have been utilized in traditional

medicine as folkloric medicine to control diabetes. *Juniperus* (*Cupressaceae*: family) is one of these plants. It is a genus of around 75 species of ever green trees found worldwide (**Raina et al., 2019**). *Juniperus phoenicea* (Arar is a local name), is a plant that grows up to around eight m in height and is extensively dispersed in the south of Saudi Arabia and Mediterranean region (**Raina et al., 2019**). *J. phoenicea* fruits were discovered in old Egyptian tombs and were grown in the Mediterranean area and Sinai. A phytochemicals analysis of the plant's fruits and leaves revealed that they have a high carbohydrate content, and/or essential oil, glycosides, flavonoids, sterols and/or triterpenes (**Al Masoudi et al., 2023**). The *J. phoenicea* species' leaves are applied in conventional medicine to treat a variety of diseases that include diabetes, bronchopulmonary disorders, and as a therapy for diuresis (**Barnawi et al., 2021**). *J. phoenicea* has also been found to have antimicrobial, antioxidant, and hepatoprotective characteristics (**Laouar et al., 2017**). Several studies have shown that *J. phoenicea* cultivated in various geographical zones can reduce the development of certain cancer cells (**Al Groshi et al., 2018**).

Origins in Europe, Mediterranean area, Central Asia, and , and North Africa are hawthorn or *Crataegus* species. They go by the name "Hawthorn" as well. Most species of hawthorn have straight, long, and pointed thorns. According to **Bouaziz et al. (2016)**, *C. azarolus* is the predominant plants in the Mediterranean area. Recent research reveals that major *crataegus* species offer advantageous impacts on human well-being. Extracts obtained from the plant's aboveground sections exhibit various biological actions, including anti-inflammatory, vasorelaxant, antihyperglycemic, and hypolipidemic (**Wang et al., 2019**). Moreover, experts have stated that the *C. azarolus* leaves have a high concentration of phenolic substances and have considerable antibacterial and antioxidant properties (**Yahyaoui et al., 2019**). Evidence indicates that the consumption of the fruits and leaves extracts of *Crataegus* is safe for humans (**Triki et al., 2022**). Bioactive substances such as flavonoids and polyphenols, including hyperoside, quercetin, and isoquercitrin, have been discovered in *C. azarolus* (**Rababa'h et al., 2020**).

Therefore this study aimed to examine how the alcohol extracts of *J. phoenicea* and *C. azarolus* leaves affect the anti-hyperglycemic in

addition to anti-hyperlipidemic capabilities of rats with streptozotocin-induced diabetes.

## **Materials and Methods**

### **Plant Material:**

Dehydrated Dry leaves of *Juniperus Phoenicea* and *Crataegus azarolus* were locally obtained from Egypt (from Haraz Shop in Cairo for medicinal plants and herbs).

### **Rats:**

Thirty rats (male albino) (*Sprague Dawely* strain) weighing  $200\pm 10$  g were acquired from the animals house in National Research Center, Giza, Egypt.

### **Streptozotocin and Biochemicals Kit:**

El-Gomhoryia Company for Chemicals in Cairo, Egypt, supplied the streptozotocin (STZ). Gamma Trading Corporation in Egypt provided kits for evaluating the serum lipid profile, radioimmunoassay kits for insulin hormones , and glucose enzymatic kits to calculating blood glucose (BG), and. Gama Trading Company, Dokki, Egypt, provided the other biochemical kits.

### **Measurment of the chemical composition of herbs:**

Chemical composition (i.e. protein, carbohydrate, fibre, fat, and ash) was assessed according to (AOAC , 2000).

### **Preparation of alcohol herbal extracts:**

To create the extract, 400 g of powdered of dried leaves from *J. phoenicea* and *C. azarolus* were soaked in two L of 90% ethanol and refrigerated while being shaken daily for five days. Afterward, the solution was percolated five to seven times until completely depleted. The resulting ethanolic extracts were then concentrated by a vacuum rotatory evaporator at a temperature of  $50^{\circ}\text{C}$  (Shalaby and Hamowieh, 2010).

### **Diabetes Induction:**

Rats were induced with diabetes through a single intraperitoneal injection of STZ produced freshly, with a dosage of  $65\text{ mg kg}^{-1}$  body

weight. The injection was diluted in 0.02 milliliter of 0.05 Molar citrate buffer with a pH of 4.5, as mentioned in **Nafiu et al. (2011)**. After a period of 72 hours, blood samples were taken from each rat's orbital plexus vein using tiny capillary tubes to verify the occurring of diabetes by testing the level of blood glucose. A diagnosis of diabetes was made once non-fasting blood glucose concentrations reached  $200 \text{ mg dL}^{-1}$ , according to **Al-Malki and El Rabey (2015)**.

### **Experimentation and dividing of rats:**

Thirty-six male adult rats (Sprague-Dawley) weighing  $200 \pm 10 \text{ g}$  were accommodated in an airy environment. The rats were kept to adapt for a week before starting the experiment, and they were fed a basal diet as outlined by **Reeves et al. (1993)**. The rats were then separated into six groups (each comprising of six rats). Group 1 acted as a normal control and received a baseline diet. Diabetes was stimulated in the other five groups by an intravenous STZ injection. Using a specialized approach, blood glucose concentrations were monitored starting on the third day after STZ injection, and continued until diabetes was confirmed (defined as blood glucose concentrations  $> 200 \text{ mg dl}^{-1}$ ) (**Al-Malki and El Rabey, 2015**). The untreated rats with diabetes comprised group (2) (as a (+) positive control), while group 3 and group 4 comprised of diabetic rats, that were fed a standard diet, and given oral doses of *J. phoenicea* extract at  $200 \text{ mg kg}^{-1} \text{ b.wt day}^{-1}$  and  $400 \text{ mg kg}^{-1} \text{ b.wt day}^{-1}$ , respectively, for a duration of four weeks (**Al-Ahdab, 2017**). On the other hand, **group 5 and group 6** were given the extract of *C. azarolus* in the mouse for four weeks in dosages of  $200$  and  $400 \text{ mg kg}^{-1} \text{ b.wt day}^{-1}$  respectively (**Pirmoghani et al., 2019**). Weekly, the animals' body weight and feed intake were recorded. Following the experimental time, the animals were weighed and then subjected to an overnight fast and sacrificed under the effects of mild ether anaesthesia.

### **Biological evaluation**

After four weeks, the experiment was finished and body weight growth, feed efficiency ratio, feed intake, and relative organ weights were measured (**Chapman et al., 1959**).

### **Serum biochemical analysis:**



After rat sacrifice, a blood sample was taken from each rat's hepatic portal vein in a dried washed centrifuge tube. Serum was cautiously extracted from blood samples by centrifugation at 3500 rpm (for 15 minutes) at room temperature, moved to a clean dry ebendorf tube, and stored at - 20 °C for further analyses. Glucose enzymatic kits were used for measuring blood glucose (BG) (**Siest *et al.*, 1981**). A particular antibody radioimmunoassay kit was used to determine serum insulin levels (**Yallow and Bauman, 1983**).

Alanine aminotransferase and aspartate aminotransferase (ALT and AST) were determined (**Bergmeyer *et al.*, 1986**); serum (ALP) was colorimetricly determined by (**Roy, 1970**); albumin (**Drupt, 1974**) ; and total protein (**Sonnenwirth and Jaret, 1980**). Globulin was estimated through the next equation :

Globulin = total protein – albumin (**Busher, 1990**)

Serum uric acid was assessed by the enzymatic colorimetric technique (**Fossati *et al.*, 1980**). Serum creatinine levels were determined by the calorimetric method (**Husdan and Rapoport, 1968**). Serum urea nitrogen levels were measured in the serum (**Patton and Crouch, 1977**).

Total cholesterol, triglycerides, and HDL-C were determined in the serum (**Allain *et al.*, 1974**; **Trinder and Ann, 1969**; **Lopes- Virella *et al.*, 1977**). measured in the serum. Serum VLDL-C and LDL-c were estimated (**Friedwald *et al.*, 1972**). The atherogenic index was estimated by using the formula:  $\log (TG/HDL-C)$  (**Nwagha, 2010**).

Serum tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) was evaluated as an inflammatory indicator (**Luo *et al.*, 2014**). The serum enzymes creatine kinase (CK) lactate dehydrogenase (LDH) were estimated by (**Uotila *et al.*, (1981)**).

Malondialdehyde (MDA) (**Yoshioka *et al.*, 1979**), declined glutathione (GSH) (**Beutler *et al.*, 1963**) and superoxide dismutase (SOD) (**Giannopolitis and Ries, 1977**) were estimated in the homogenized pancreas.



## Statistical analysis:

Statistical analysis was done with the analysis of variance (one-way) (ANOVA) and Duncan assess in the SPSS software (Chicago, IL, USA). The results were presented as average  $\pm$  SD. The difference among averages was considered significant at  $P < 0.05$  (Snedecor and Cochran, 1989).

## Results

### 1. Chemical analysis of *Juniperus phoenicea* and *Crataegus azarolus*

The data in Table 1 displayed that *J. phoenicea* and *C. azarolus* chemical analysis have the highest values of carbohydrate ( $66.66\% \pm .04$ ), crude protein ( $10.61\% \pm .02$ ) and ash ( $8.69\% \pm .020$ ), while *J. phoenicea* has the highest values of fiber ( $17.10\% \pm .22$ ) and crude fat ( $5.28\% \pm .020$ ).

**Table (1): Chemical analysis of leaves of *Juniperus phoenicea* and *Crataegus azarolus* g 100g<sup>-1</sup>**

Components %	<i>Juniperus phoenicea</i>	<i>Crataegus azarolus</i>
Carbohydrate	$64.80 \pm .05^b$	$66.66 \pm .04^a$
Fiber	$17.10 \pm .22^a$	$13.27 \pm .02^b$
Crude fat	$5.28 \pm .020^a$	$0.76 \pm .02^b$
Crude protein	$5.78 \pm .02^b$	$10.61 \pm .02^a$
Ash	$7.04 \pm .04^b$	$8.69 \pm .020^a$

At significance level of ( $P < 0.05$ ), the letters in the same column with different letters are statistically significant.

### 2. Biological evaluation

The results in Table 2 display that the average values of FI, BWG %, FER and relative pancreas weight in the + control group were lower significantly than the - control group. Conversely, all measurements in the remedy groups were increased significantly at ( $P < 0.05$ ). Therefore, the best relative pancreas mass and FI results were recorded at 400 mg kg<sup>-1</sup> BW *J. phoenicea*. Simultaneously, the preferable BWG % and FER findings were noticed in *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW).

**Table (2): The effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on feed intake (FI), body weight gain (BWG %), feed efficiency ratio (FER) and relative pancreas mass in diabetic rats (M±SD)**

Parameters	FI (g)	BWG (%)	FER(g)	Pancreas (%) Row
Groups	28days			
Control -ve	500.13 ±5.20 <sup>a</sup>	40.38±3.11 <sup>a</sup>	0.122±0.004 <sup>a</sup>	0.60±0.03 <sup>a</sup>
Control +ve	378.40±2.23 <sup>f</sup>	13.80±2.05 <sup>d</sup>	0.053±0.005 <sup>e</sup>	0.36±0.04 <sup>d</sup>
<i>J. phoenicea</i> 200 mg	423.33 ±2.81 <sup>d</sup>	24.53±1.88 <sup>c</sup>	0.085±0.002 <sup>c</sup>	0.49±0.01 <sup>bc</sup>
<i>J. phoenicea</i> 400 mg	464.67±3.84 <sup>b</sup>	35.17±3.06 <sup>b</sup>	0.112±0.008 <sup>b</sup>	0.52±0.02 <sup>b</sup>
<i>C. azarolus</i> 200 mg	414.60± 40 <sup>e</sup>	20.82±4.01 <sup>c</sup>	0.075 ±0.006 <sup>d</sup>	0.46±0.02 <sup>c</sup>
<i>C. azarolus</i> 400 mg	442.93±2.97 <sup>c</sup>	31.45±1.78 <sup>b</sup>	0.104 ±0.005 <sup>b</sup>	0.50±0.02 <sup>bc</sup>

At significance level of ( $P < 0.05$ ), the letters in the same column with different letters are statistically significant.

### 3. Blood glucose and serum insulin

According to data in Tables (3), the (+ve) control group's average values of blood glucose were significantly higher than the (-ve) control. All handled groups recorded a significant decline in all parameters than the control (+ve). The best results were recorded in rats receiving *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW), while serum insulin in the positive control was significantly decreased than the normal rats. The highest serum insulin concentration was recorded in treated groups with *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW).

**Table (3): The effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on blood glucose and serum insulin in diabetic rats (M±SD)**

Parameters Groups	Blood glucose (mg dl <sup>-1</sup> )	Serum insulin (pg ml <sup>-1</sup> )
Control -ve	80.0±2.0 <sup>e</sup>	287.50±4.78 <sup>a</sup>
Control +ve	403.67±3.51 <sup>a</sup>	81.00±3.0 <sup>e</sup>
<i>J. phoenicea</i> 200 mg	271.33±3.05 <sup>c</sup>	152.33±4.517 <sup>c</sup>
<i>J. phoenicea</i> 400 mg	124.67±4.04 <sup>d</sup>	256.83±6.53 <sup>b</sup>
<i>C. azarolus</i> 200 mg	281.67±3.05 <sup>b</sup>	142.17±5.01 <sup>d</sup>
<i>C. azarolus</i> 400 mg	128.67±3.21 <sup>d</sup>	249.00±6.00 <sup>b</sup>

At significance level of ( $P < 0.05$ ), the letters in the same column with different letters are statistically significant.

#### 4. Liver functions

The data in Table 4 demonstrate that the average value of AST, ALT and ALP levels in the +ve group was significantly higher compared with the -ve group. However, all other examined groups decreased significantly than the +ve group. The minimum AST, ALT and ALP levels were noticed in the treated group with *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW).

**Table (4): Effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on liver enzyme (AST, ALT and ALP) in diabetic rats (M±SD)**

Parameters Groups	AST (U L <sup>-1</sup> )	ALT (U L <sup>-1</sup> )	ALP (U L <sup>-1</sup> )
Control -ve	91.0± 4.0 <sup>e</sup>	29.0±3.0 <sup>d</sup>	251.33±4.51 <sup>d</sup>
Control +ve	200.33±4.51 <sup>a</sup>	69.0±2.0 <sup>a</sup>	463.0 ±7.0 <sup>a</sup>
<i>J. phoenicea</i> 200 mg	180.0±3.0 <sup>c</sup>	52.0±3.0 <sup>b</sup>	388.33±4.51 <sup>b</sup>
<i>J. phoenicea</i> 400 mg	112.67±2.31 <sup>d</sup>	35.0±3.0 <sup>c</sup>	277.67±2.51 <sup>c</sup>
<i>C. azarolus</i> 200 mg	191.0±4.0 <sup>b</sup>	56.0±4.0 <sup>b</sup>	390.0±5.0 <sup>b</sup>
<i>C. azarolus</i> 400 mg	116.0±4.0 <sup>d</sup>	38.0±2.0 <sup>c</sup>	284.0±3.61 <sup>c</sup>

At significance level of ( $P < 0.05$ ), the letters in the same column with different letters are statistically significant.

## 5. Serum albumin, globulin and total protein

The obtained numbers in Table 5 show that the average values of total protein, albumin, and globulin in the +ve control declined significantly than the -ve one. Conversely, the other investigated groups showed a significant increase than the +ve control. The highest total protein and albumin were noted in the handled group with *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW). However, the maximum globulin was recorded in group *J. phoenicea* (200 mg kg<sup>-1</sup> BW).

**Table (5): Effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on total protein, albumin and globulin in diabetic rats (M±SD)**

Parameters Groups	T. Protein (g dl <sup>-1</sup> )	Albumin (g dl <sup>-1</sup> )	Globulin (g dl <sup>-1</sup> )
<b>Control -ve</b>	8.40±0.01 <sup>a</sup>	4.64±0.04 <sup>a</sup>	3.76±0.04 <sup>a</sup>
<b>Control +ve</b>	5.93±0.03 <sup>d</sup>	3.06±0.03 <sup>d</sup>	2.87±0.03 <sup>d</sup>
<b><i>J. phoenicea</i> 200 mg</b>	7.17±0.03 <sup>c</sup>	3.60±0.06 <sup>c</sup>	3.57±0.03 <sup>b</sup>
<b><i>J. phoenicea</i> 400 mg</b>	7.91±0.04 <sup>b</sup>	4.38±0.02 <sup>b</sup>	3.53±0.03 <sup>bc</sup>
<b><i>C. azarolus</i> 200 mg</b>	7.12±0.02 <sup>c</sup>	3.59±0.03 <sup>c</sup>	3.53±0.03 <sup>bc</sup>
<b><i>C. azarolus</i> 400 mg</b>	7.88±0.03 <sup>b</sup>	4.42±0.08 <sup>b</sup>	3.46±0.07 <sup>c</sup>

At significance level of ( $P < 0.05$ ), the letters in the same column with different letters are statistically significant.

## 6. Heart enzyme

Table 6 displays the serum LDH and CPK levels that were increased in the unhealthy rats than in the standard control. It was lower significantly in all treatments than the +ve control. The most effective outcomes in LDH were noted with *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW). In comparison, a preferable finding in CPK was found in *J. phoenicea* (400 mg kg<sup>-1</sup> BW).

**Table (6): Effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on heart enzyme (LDH and CPK) in diabetic rats (M±SD)**

Parameters	LDH (U L <sup>-1</sup> )	CPK (U L <sup>-1</sup> )
<b>Groups</b>		
Control –ve	450.0 ±10.0 <sup>e</sup>	350.0 ± 5.0 <sup>f</sup>
Control +ve	717.0 ±17.0 <sup>a</sup>	670.0 ± 12.0 <sup>a</sup>
<i>J. phoenicea</i> 200 mg	600.0 ±10.0 <sup>c</sup>	520.0 ± 9.0 <sup>c</sup>
<i>J. phoenicea</i> 400 mg	500.0 ±12.0 <sup>d</sup>	400.0 ±10.0 <sup>e</sup>
<i>C. azarolus</i> 200 mg	630.0 ±13.0 <sup>b</sup>	550.0 ±15.0 <sup>b</sup>
<i>C. azarolus</i> 400 mg	516.67 ± 6.66 <sup>d</sup>	500.0 ±11.0 <sup>d</sup>

At significance level of ( $P < 0.05$ ), the letters in the same column with different letters are statistically significant.

## 7. Kidney functions

The obtained results in Table 7 show that + ve control group's average serum creatinine, urea, and uric acid levels were higher significantly than their counterparts in the – ve negative control. Compared to the + ve control, all parameters in the remedy categories reduced significantly ( $P < 0.05$ ). The best creatinine and urea findings were found in *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW). At the same time, the preferable serum uric acid finding was found in *J. phoenicea* (400 mg kg<sup>-1</sup> BW).

**Table (7): Effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on creatinine, urea and uric acid in diabetic rats (M±SD)**

Parameters	Creatinine (mg dl <sup>-1</sup> )	Urea (mg dl <sup>-1</sup> )	Uric acid (mg dl <sup>-1</sup> )
Groups			
<b>Control -ve</b>	0.43±0.03 <sup>d</sup>	14.33± 2.97 <sup>d</sup>	1.95± 0.15 <sup>d</sup>
<b>Control +ve</b>	1.55±0.05 <sup>a</sup>	81.67± 3.48 <sup>a</sup>	3.74 ± 0.04 <sup>a</sup>
<b><i>J. phoenicea</i> 200 mg</b>	0.99±0.04 <sup>b</sup>	50.43± 4.76 <sup>b</sup>	3.04± 0.21 <sup>b</sup>
<b><i>J. phoenicea</i> 400 mg</b>	0.60±0.02 <sup>c</sup>	28.20± 3.95 <sup>c</sup>	2.29± 0.26 <sup>cd</sup>
<b><i>C. azarolus</i> 200 mg</b>	0.98±0.04 <sup>b</sup>	55.63±4.20 <sup>b</sup>	3.08± 0.20 <sup>b</sup>
<b><i>C. azarolus</i> 400 mg</b>	0.61±0.03 <sup>c</sup>	29.53±3.87 <sup>c</sup>	2.34± 0. 26 <sup>c</sup>

At significance level of ( $P < 0.05$ ), the letters in the same column with different letters are statistically significant.

### 8. Lipid profile

The data in Tables 8 and 9 reveal that the +ve control average values of the lipid profile (TC, LDL, AI, TG, and VLDL) increased significantly when than the - ve control group. All treatments recorded a significant decline in all parameters than the +ve control. The best data was noticed in the rats that received *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW) in AI, TG and VLDL. While the best results in TC and LDL were found in rats receiving *J. phoenicea* (400 mg kg<sup>-1</sup> BW). On the other hand, HDL in the +ve control was significantly decreased than the normal rats. The highest HDL content was recorded in the group that treated with *J. phoenicea* (400 mg kg<sup>-1</sup> BW).

**Table (8): Effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on (TC, HDL, LDL and AI) in diabetic rats (M±SD)**

Parameters	TC	HDL	LDL	AI
Groups	(mg dl <sup>-1</sup> )	(mg dl <sup>-1</sup> )	(mg dl <sup>-1</sup> )	(mg dl <sup>-1</sup> )
<b>Control -ve</b>	130.0±4.0 <sup>f</sup>	40.0±4.00 <sup>a</sup>	75.53±4.00 <sup>e</sup>	1.82±0.16 <sup>c</sup>
<b>Control +ve</b>	200.0±5.0 <sup>a</sup>	20.0±3.00 <sup>d</sup>	153.40±3.64 <sup>a</sup>	6.76±1.11 <sup>a</sup>
<b><i>J. phoenicea</i> 200 mg</b>	179.67±4.51 <sup>c</sup>	25.33±2.51 <sup>c</sup>	132.73±1.81 <sup>b</sup>	4.29±0.39 <sup>b</sup>
<b><i>J. phoenicea</i> 400 mg</b>	149.67±4.51 <sup>e</sup>	37.00±3.00 <sup>ab</sup>	95.47±1.97 <sup>d</sup>	2.34±0.25 <sup>c</sup>
<b><i>C. azarolus</i> 200 mg</b>	189.0±4.0 <sup>b</sup>	26.0±2.65 <sup>c</sup>	141.40±7.15 <sup>b</sup>	4.17±0.30 <sup>b</sup>
<b><i>C. azarolus</i> 400 mg</b>	167.0±3.0 <sup>d</sup>	32.0±2.00 <sup>b</sup>	117.53±0.50 <sup>c</sup>	2.73±0.09 <sup>c</sup>

At significance level of ( $P < 0.05$ ), the letters in the same column with different letters are statistically significant.

**Table (9): Effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on (TG and VLDL ) in diabetic rats (M±SD)**

Parameters	TG (mg dl <sup>-1</sup> )	VLDL (mg dl <sup>-1</sup> )
Groups		
<b>Control -ve</b>	72.33±3.51 <sup>d</sup>	14.47±0.70 <sup>d</sup>
<b>Control +ve</b>	133.0±4.00 <sup>a</sup>	26.60±0.80 <sup>a</sup>
<b><i>J. phoenicea</i> 200 mg</b>	108.0±2.00 <sup>b</sup>	21.60±0.40 <sup>b</sup>
<b><i>J. phoenicea</i> 400 mg</b>	86.0±3.00 <sup>c</sup>	17.20±0.60 <sup>c</sup>
<b><i>C. azarolus</i> 200 mg</b>	108.0 ±3.00 <sup>b</sup>	21.60±0.60 <sup>b</sup>
<b><i>C. azarolus</i> 400 mg</b>	87.33±2.52 <sup>c</sup>	17.47±0.50 <sup>c</sup>

Atsignificance level of ( $P < 0.05$ ), the letters in the same column with different letters are statistically significant.



## 9. Serum tumor necrosis factor - $\alpha$ (TNF- $\alpha$ )

Table 10 shows that the average values of serum TNF-  $\alpha$  were higher in the unhealthy group than in the standard control rats. It was lower significantly in all treatment groups than the +ve control. The most effective outcomes were noted with *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW).

**Table (10): Effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on serum tumor necrosis factor - $\alpha$  (TNF-  $\alpha$ ) in diabetic rats (M $\pm$ SD)**

Parameters	TNF- $\alpha$ (pg ml <sup>-1</sup> )	
Groups		
Control -ve	21.47 $\pm$ 1.72 <sup>e</sup>	
Control +ve	60.85 $\pm$ 2.32 <sup>a</sup>	
<i>J. phoenicea</i> 200 mg	40.25 $\pm$ 1.05 <sup>c</sup>	significance 0.05), the same different statistically
<i>J. phoenicea</i> 400 mg	26.57 $\pm$ 1.60 <sup>d</sup>	
<i>C. azarolus</i> 200 mg	44.03 $\pm$ 1.74 <sup>b</sup>	
<i>C. azarolus</i> 400 mg	27.42 $\pm$ 2.73 <sup>d</sup>	

At level of ( $P <$  letters in the column with letters are significant.

### 10.

## Antioxidant enzymes (GSH, SOD), malondialdehyde (MDA) in pancreas tissue.

Table 11 illustrates that the activities of pancreatic reduced glutathione (GSH) and superoxide dismutase (SOD) were lowered significantly in the +ve positive control group than the -ve one. At the same time, they rose in other groups compared with the +ve control rats. The best findings in GSH and SOD were noticed in groups of *J. phoenicea* and *C. azarolus* (400 mg kg<sup>-1</sup> BW). The average value of MDA was significantly higher in the +ve control than in the -ve one; however, their values were significantly lower in other groups than in the +ve control. The best finding was got in *J. phoenicea* (400 mg kg<sup>-1</sup> BW).

**Table (11): Effect of *Juniperus phoenicea* and *Crataegus azarolus* alcohol extracts on Antioxidant enzymes (GSH, SOD), malondialdehyde (MDA) in diabetic rats (M±SD)**At significance level of ( $P < 0.05$ ), the letters in the same column with different

Parameters	GSH	SOD	MDA
Groups	(mmol g <sup>-1</sup> )	(U g <sup>-1</sup> )	(nmol g <sup>-1</sup> )
<b>Control -ve</b>	2.06± 0.16 <sup>a</sup>	133.38± 3.19 <sup>a</sup>	6.12±1.21 <sup>d</sup>
<b>Control +ve</b>	0.51± 0.03 <sup>c</sup>	60.75 ± 1.98 <sup>d</sup>	27.59±2.71 <sup>a</sup>
<b><i>J. phoenicea</i> 200 mg</b>	1.14± 0.17 <sup>b</sup>	92.58± 2.86 <sup>c</sup>	15.06±1.15 <sup>b</sup>
<b><i>J. phoenicea</i> 400 mg</b>	1.89± 0.23 <sup>a</sup>	127.30± 3.20 <sup>b</sup>	8.19±2.05 <sup>cd</sup>
<b><i>C. azarolus</i> 200 mg</b>	1.06±0.17 <sup>b</sup>	87.55± 3.25 <sup>c</sup>	15.77±1.39 <sup>b</sup>
<b><i>C. azarolus</i> 400 mg</b>	1.85±0.23 <sup>a</sup>	126.33± 3. 91 <sup>b</sup>	10.0±2.0 <sup>c</sup>

letters are statistically significant.

## Discussion

Diabetes is intimately linked to oxidative stress, which can harm both islet  $\beta$  cells and insulin signalling pathways, possibly leading to catastrophic effects such as cardiovascular disease and nephropathy (Zhang *et al.*, 2019). According to recent research, several medicinal herbs are recognized as valuable bases for conventional medications, and many contemporary medications are derived from these plants (Dar *et al.*, 2019). Though, the use of herbs in remedy is mostly relied on their physiologically active chemicals, which offer a diversity of therapeutic capabilities such as antioxidant, anti-diabetic, anti-inflammatory, and anti-hypercholesterolemic properties. (Neeta *et al.*, 2015). As a consequence, many researches have been found to evaluate plant secrets. Thus, the current research was designed to assess the possible anti-diabetic benefits of alcohol extract of *J. phoenicea* and *C. azarolus*. Özcan *et al.* (2005)

found that the chemical composition of hawthorn (*Crataegus* spp.) was 34.02 kcal g<sup>-1</sup> energy, 2.48% protein, 0.87% oil, 4.67% cellulose, and 2.28% ash. **Chen et al. (2019)** revealed that protein level was 0.83% for *C. azarolus* pulps (CAP) and 5.68% for seeds (CAS). The CAP had 82.35% of carbohydrate and 52.86% neutral sugar level, respectively, that were more than those of the CAS; 64.93% of carbohydrate and 45.25% of neutral sugar. **Zhang et al., (2022)** stated that the chemical composition of hawthorn (*C. azarolus*) was recorded as 7 mg g<sup>-1</sup> of protein, 2 mg g<sup>-1</sup> of fat, and 68.98% of water.

In this work, rats that STZ-induced diabetic exhibited a decline in BWG%, and FER when than the-ve group. The current results were accorded with **(Suryanarayana et al., 2005)** and **(Gupta et al., 2012)** where they stated that STZ-induced diabetic rats displayed weight loss than rats not treated with STZ. **Kota et al. (2012)** stated that an relationship between hyperglycemia and loss of body weight in diabetic rats, diabetes-induced weight loss, as well as the body's inability to retain or utilize glucose, promote hunger and weight loss. Body weight loss in STZ diabetic rats is ascribed to an increment in blood glucose with insulin inhibition, a reduction in tissue proteins, and an increment in muscle wasting **(Zafar and Naqvi, 2010)**. These obtained data were in accordance with **Yang and Hyung-Sub, (2018)** where they show that the significant loss in body weight in sSTZ-induced diabetic rats is attributable to relative or absolute insulin insufficiency or reduced protein creation in the tissues **(Sadri et al., 2017)**.

The findings of the ongoing work also accord with **Tebboub and Kechrid (2021)** and **Triki et al. (2022)** where they noticed that the group of individuals with diabetes displayed a noteworthy decline in body weight contrary to the healthy group. This outcome can be attributed to a disruption in their metabolic state, suggesting that the inability to utilize carbohydrates as a source of energy and the elevated breakdown of fat and protein mass may be the causes of weight loss. In this regard, **Gadewar et al. (2023)** discovered that significant weight loss is one of the most prominent aspects of STZ-induced diabetes, because of greater protein waste and the non-availability of glucose for energy use. The present results were supported by **Saswata et al. (2013)** proposed that the rise in body weight was gradually seen in STZ diabetic animals that treated with

the extract of *J. Phoenicea* leaves might be attributed to the preserved contents of glucose and insulin due to the antioxidant actions of the extract of *J. Phoenicea* leaves. **Keskes et al. (2014)** discovered that *J. Phoenicea* leaves extract enhanced the feed intake and body weight in diabetic rats given *J. Phoenicea* leaves extract compared to diabetic untreated rats because of its potential to prevent angiogenesis in adipose tissues and diminish preadipocyte variation .

**Dall et al. (2016)** underlined that the streptozotocin (STZ) group lost body weight whereas the hawthorn leaf (*C. azarolus*) flavonoids group increased body weight since hawthorn leaf flavonoids had a preventive impact on diabetic cardiomyopathy rats to some extent. **Triki et al. (2022)** observed that *C. azarolus* improved body weight in diabetic rats. The considerable increment in animal body weight might be attributed to increased food intake and protein synthesis. As well, *C. Azarolus* has also been shown to be capable of reversing gluconeogenesis and controlling protein loss. **Kumar et al. (2012)** found a similar conclusion, indicating that streptozotocin injection resulted in a significant increase in plasma glucose and a drop in insulin level because STZ caused the death of  $\beta$ -cells in the islets of Langerhans, leading to in cell degranulation and a decrease in insulin production as suggested by **Zhang and Tan, (2002)**. Our study goes along the same lines as **Sekkin et al. (2015)** who noticed that an injection for one time of STZ causes the advance of DM, which is featured by high blood glucose and reduced serum insulin levels. This happened due to STZ-induced islet cell damage,  $\beta$  cell death within 24 hours, and the activation of an inflammatory process that resulted in macrophage and lymphocyte infiltration. STZ-induced oxidative damage in rats is directly linked to chronic inflammatory disease, which may lead to tissue damage **Xu et al. (2014a)**.

These conclusions are accorded with **El-Sawi et al. (2015)**, who evaluated the influences of *J. phoenicea* fruits and leaves on diabetic animals. The obtained data demonstrated that *J. phoenicea* extracts reduced blood glucose levels. Animal DM inductions caused injury to pancreatic tissues. This damage can cause histopathological changes in pancreatic islets, in addition to a rise in blood glucose as well a drop in insulin levels (**Wahba et al., 2016, Elkotby et al., 2018**). Extraction of *J. Phoenicea* leaf decreases fasting blood glucose concentration in a dose-

dependent method, most likely via boosting insulin production from remaining beta cells (**Karuppusamy and Thangaraj, 2012**). In accordance with the present findings, which were supported by **Al-Ahdab (2017)**, the *J. phoenicea* extract lowered blood glucose and rised insulin concentrations, normalized serum contents of liver enzymes and biochemical markers of kidney functions, and improved the lipid profile in STZ rats treated when comparing with untreated animals.

These data also are in accordance with **Goyal et al. (2016)**, who discovered that the leaf flavonoids of hawthorn significantly lowered blood glucose levels when compared to diabetic cardiomyopathy rats due to hawthorn leaf flavonoids' preventive impact. **Abu-Gharbieh and Shehab (2017)** reported that treating diabetic rats with the extraction of *C. azarolus* leaves resulted in a reduced baseline blood glucose concentration. According to **Abu-Gharbieh and Shehab (2017)**, oral administration of *C. azarolus* demonstrated strong antihyperglycemic influence and noticeable performance in glucose tolerance (OGTT), confirming the efficiency of *C. azarolus* extract as a hypoglycemic agent by postponing the ingestion of carbohydrates and reducing blood glucose concentration. The current study's results correspond with those of **Zafar et al. (2009)**; **Najla et al., (2012)**; and **Soliman, (2013)**, who indicated that STZ diabetes produces increases in amino transferases as well as a rise in blood ALP activity. The increased oxidative stress caused by STZ diabetes may be the reason for the hepatotoxicity that occurs when ALT and AST leak from liver cells (**Okechukwu et al. 2013**). In this regard, **Ramesh et al. (2010)** and **Dewanjee et al. (2020)** stated that in hyperglycemia-related liver disorders, oxidative stress is critical. STZ-induced necrosis raises blood ALT and AST levels due to cytosol leak from the liver, and excessive contents of these enzymes imply possible liver injury and hepatotoxicity. These findings were in accordance with the results of **Makena et al. (2018)**, who found that ALT and AST levels were higher in the diabetic rats than the control rats. In Diabetes Mellitus, ALT and AST are linked to the transformation of amino acids into keto acids.

These results also were verified by **Yang et al. (2011)** who found that *J. phoenicea* berries have elevated plasma total protein and albumin levels indicating hepatoprotective activity. Protein production stimulation

has been proposed as a hepatoprotective process that speeds the self-renewal and formation of liver cells. **Mistry et al., (2013)** stated that administration of *J. phoenicea* berries has significantly protected hepatocytes injury boosted by CCl<sub>4</sub> through decreasing AST and ALT levels, which is a sign of the hepatic cell renewal process. However, a decrease in ALP levels with contemporaneous depletion shows that biliary function is stable. The significant reducing in blood liver enzymes level (ALT, AST, and ALP) in diabetic animals demonstrated the hepatoprotective activity of the extract of *J. Phoenicea* Leaves described in this work; this result was consistent with its polyphenols (**Salma et al., 2017**). **Dessoky et al., (2020)** also found that gamma-irradiated rats that pre-treated the extraction of *J. phoenicea* leaves had a significant rise in total protein, albumin, and globulin concentrations and a significant decrease in liver enzymes and lipid profile indicators compared to healthy rats because *J. phoenicea* leaves could be a possible source of therapeutic antioxidants. This result was also supported by **Nasir et al. (2013)** who noted that *C. azarolus* methanolic extract administration stabilizes the endoplasmic reticulum, leading to a rise in total protein levels and protein production in the liver. Moreover, **Othman and Mustafa (2017)** observed that hawthorn dramatically lowers ALP levels while increasing SGOT and SGPT levels when compared to a high-triglyceride diet.

This finding was confirmed by **Mohammed et al. (2016)** and **Suanarunsawat et al. (2016)** who stated that the CK and LDH levels were significantly higher in diabetic animals because the integrity of cardiac tissues' cellular membranes may be disturbed. **Mandlem and Annapurn, (2017)** found that increased levels of serum CK and LDH in STZ diabetic animals may be due to their increased produce from cardiac necrotic tissues. As well, fibrosis and necrosis of cardiac muscle fibers, changes in oxidative stress, and cardiac biomarker indicators were noticed in diabetic animals and patients.

The results also accorded with those of **Cui et al. (2006)** and **Ali and Shapour (2013)** who showed that rats treated with hawthorn (*C. azarolus*) induced decreased cardiac enzymes (creatinine kinase and lactate dehydrogenase) suggesting that hawthorn may be moderating the harmful impact of Doxorubicin as a result of hawthorn's antioxidant scavenging potential.



According to **Xu et al. (2014 b)** and **Guo et al. (2021)**, STZ-induced diabetic nephritis group rats had higher levels of BUN and uric acid than the untreated and treated groups. Kidney disease progresses through modifying kidney hemodynamics, resulting in glomerulosclerosis, kidney failure, and proteinuria. With diabetic nephropathy, a nitrogen imbalance combined with decreased protein synthesis causes the production of non-protein nitrogenous substances such as creatinine and BUN. The untreated diabetic group showed significantly higher blood levels of BUN, UA, and Cr when than normal control group .These data is in accordance with **Ana et al. (2009)** who signified that enhanced kidney activities are indications of kidney disorder in diabetic patients, but after therapies with extract of *J. Phoenicea* leaves, a significant decrease was noticed in comparison to the + ve group, which may be due to the antioxidant activities. These obtained results were also accorded with **Mall et al. (2016)** where they stated that the antioxidant and hypoglycemic of *J. Phoenicea* leaves extract reduced oxidative stress, leading to reduce the serum content of BUN, UA, and Cr.

This conclusion is reinforced by **Qin et al. (2019)** who observed that Hawthorn (*Crataegus azarolus*) significantly lowered blood urea nitrogen in DKD rats since hawthorne may prevent kidney damage and improve kidney function. This might be done by modulating p38MAPK activation and reducing oxidative stress damage.

The results are also reinforced by **Kusunoki et al. (2000)** who described that hyperlipidemia is a result of elevated lipolysis, which results in an increase in free fatty acids due to s and glycerol, which the liver uses to create acetyl Co A. Because of the rise in intestine acyl coenzyme A, acetyl Co A is a precursor for cholesterol formation in STZ-induced diabetic rats. **Zhang et al. (2016)** showed that diabetic rats had considerably higher triglycerides, cholesterol, VLDL-C, and LDL-C, and lower HDL-C. **Pirmoghani et al. (2019)** noted that the contents of TG and cholesterol in the diabetic group rose when comparing with the - ve control. Furthermore, these data are in the line with those of **Chen et al. (2020)** who found that generally diabetes is related to irregular lipid metabolism, that is a great risk reason for diabetic cardiovascular disease. Elevated concentrations of cholesterol and triglycerides were undoubtedly caused by changes in lipid metabolism under diabetes



circumstances, which eventually led to suppression of the activity of lipoprotein lipase due to insulin insufficiency and insulin resistance. These results were well-matched with **Abbas and Ali, (2021)** found that there were a significant increase in triglyceride (TG), cholesterol, (LDL), combined with significant decline in high density lipoprotein (HDL) in diabetic groups than the control groups. In this regard, **Gadewar et al., (2023)** noticed an increment in the level of serum total cholesterol and triglycerides with a reduction in the HDL level that was noticed in STZ-induced diabetic groups.

The results in agreement with those observed by **Ono et al. (2006)** revealed that treatment by *J. Phoenixea* leaves extract may inhibit pancreatic lipase activity and inhibit or delay lipid absorption. **Saswata et al. (2013)** informed that treating of diabetic animals with the extraction of *J. Phoenixea* leaves showed extremely improved consequences; there was a significance in enhancement TC, TG, LDL-C, and HDL-C concentrations than the untreated diabetic rats. **Henda et al. (2014)** found that the inhibitive powers of *J. phoenixea* extract contrary to lipase activity may be entirely coincidental with their total phenolic components.

The present findings were verified by **Jia et al. (2011)** who reported a significant decrease in the TG, TC, LDL-C, and VLDL-C levels of the group treated with the extraction of ethanolic leaves of *C. azarolus*. As well, the serum level of HDL-C was greater than its counterpart in the control rats decreasing the “Atherogenic Index” and the ratios of LDL-C: HDL-C by enhancing fatty acid absorption and oxidation while blocking fatty acid synthesis. **Pirmoghani et al. (2019)** showed that *C. azarolus* extract lowered LDL and cholesterol and rised HDL in diabetic groups.

Numerous studies have demonstrated that an increase in TNF- is connected with diabetes and insulin resistance occurrence in type 2 diabetic individuals, either directly or indirectly (**Navarro and Mora, 2006**) and (**Alexandraki et al., 2006**).

The findings are accorded with th **Mardiaha et al., (2015)**, where they demonstrated that diabetic animals had the maximum level of TNF- $\alpha$ . This was related to the inflammatory response of TNF- $\alpha$ , which resulted in  $\beta$ -cell death. A high concentration of inflammatory cells may

cause  $\beta$ -cell apoptosis, which may result in a decrease in the cell numbers of insulin-producing or a fall in insulin levels. **Elsawy and Emara (2016)** found that elevated MDA and TNF- $\alpha$  levels in diabetes mellitus were caused by increased oxidative stress or a reduction in antioxidant defence mechanisms. According to **Zhao et al. (2016)**, *J. sabina* considerably reduced the concentrations of TNF- $\alpha$ , hence *J. sabina* handled rats exhibited a significant decrease in arthritic scoring than the control group. The current results were supported by **Mostafa et al. (2018)** who noticed that *Crataegus aronia* reduced glucose concentration, serum lipids, and significantly reduced tumour necrosis factor alpha. (TNF $\alpha$ ).

The present study's results are also in the line with those of **Mahboob et al., (2005)**, **Senthilkumar et al., (2006)**, and **Gadewar et al., (2023)**, where they discovered that hyperglycemia related to diabetes produces free radicals via glucose breakdown, non-enzymatic glycation of protein, or resulting oxidative degradation, and that these free radicals induce cellular membrane destruction and lipid peroxidation in diabetes. Alterations in lipid peroxidation in diabetic rats revealed a reduction in numerous antioxidant enzymes activities, including superoxide dismutase, which have a significant function in removing the toxic compounds created during the incomplete oxidation process. Our findings suggest that hyperglycemia raises pancreatic MDA while decreasing SOD and GSH activity. These influences might be attributed to hyperglycemia's increased oxidative stress via the overproduction of ROS (**Hussein and Abu-Zinadah, 2010**). **Molehin et al. (2018)** stated that the increment in levels of ROS in diabetic animals resulted in a decline in SOD and GSH levels and an increase in MDA levels, confirming that STZ generated oxidative stress. The results were agreed with those of **Elbe et al., (2015)** and **Lo et al., (2017)** showed that a lowered catalase level and rised level of MDA in diabetic rats. The lowering in these antioxidants activity can cause an increment in hydrogen peroxide and superoxide anion in biological systems, consequently create hydroxyl radicals that result in lipid peroxidation. **Al-Attar and Alsalmi (2019)** found a decline in SOD and GSH, and significant enhancement in MDA value in diabetic rats.

*Juniperus Phoenicea* leaves extract increases pancreatic tissue MDA, GSH, and SOD levels in hyperglycemic rats. *J. Phoenicea* leaves

extract decreases oxidative stress by lowering peroxide production and enhancing antioxidant enzyme activity, according to **Barrero et al. (2004)**. **Al-Ahdab (2017)** demonstrated that *J. phoenicea* extract decreased MDA and elevated SOD and GSH concentrations in STZ diabetic rats in comparison with untreated diabetic rats. According to the current findings, which were validated by **Ayala et al., (2014)**, treatment with hawthorn leaf flavonoids boosted SOD activity and dramatically reduced the level of diabetic cardiomyopathy. Hawthorn (*C. azarolus*) leaf flavonoids may preserve diabetic cardiomyopathy rats by reducing ROS production and alleviating oxidative stress damage by antioxidant interventions. **Triki et al., (2022)** showed that treating with *C. Azarolus* resulted in a considerable decrease in the generation of malondialdehyde, an increase in GSH content, and an enhancement in SOD activity in zinc deficiency diabetic rats.

### Conclusion

Among the most crucial non-infectious diseases to affect the world in the current millennium is diabetes mellitus. It is among the most severe, long-lasting disorders of protein, lipid, and carbohydrate metabolism. Despite significant medical advancements, there is still no fully effective medication to treat diabetes mellitus. More and more research is showing that a variety of nutritious natural foods, medicinal herbs, and dietary additions have the possibility to be effective complementary therapies for the treating of type 2 diabetes and its consequences. In the current investigation, diabetic male rats were used to test the hypoglycemic effects of alcohol extracts from *Juniperus Phoenicea* and *Crataegus azarolus* leaves. Considering the outcomes of the current experiments, it can be concluded that this study demonstrates that alcoholic extracts of *J. Phoenicea* and *C. azarolus* leaves reduced the physiological alterations caused by STZ in the experimental groups. To determine the ideal dosages of such extracts as hypoglycemic drugs and to clarify their modes of action, more physiological, pharmacological, and biochemical researches are required

## References:

- Abbas, M. and Ali, M. (2021).** Protective Effects of Beetroot on Streptozotocin Induced Diabetes in Adult Male Albino Rats. *Bulletin of Egyptian Society for Physiological Sciences*, 41(2), 270-282.
- Abu-Gharbieh, E. and Shehab, N.G. (2017).** Therapeutic potentials of *Crataegus azarolus* var. eu-azarolus Maire leaves and its isolated compounds. *BMC Complementary and Alternative Medicine* 17- 218.
- Al Groshi, A., Evans, A. R., Ismail, F. M., Nahar, L. and Sarker, S. D. (2018).** Cytotoxicity of Libyan *Juniperus phoenicea* against human cancer cell lines A549, EJ138, HepG2 and MCF7. *Pharmaceutical Sciences*, 24(1), 3-7.
- Al Masoudi, L. M., Alqurashi, A. S., Abu Zaid, A. and Hamdi, H. (2023).** Characterization and Biological Studies of Synthesized Titanium Dioxide Nanoparticles from Leaf Extract of *Juniperus phoenicea* (L.) Growing in Taif Region, Saudi Arabia. *Processes*, 11(1), 272.
- Al-Ahdab, M. A. (2017).** Hypoglycemic effect of alcoholic extracts of *Pyllanthus virgatus* and *Juniperus Phoenicea* L., on streptozotocin-induced diabetic in male rats. *Life Science Journal*, 14(9) 61–70.
- Al-Attar, A. M. and Alsalmi, F. A. (2019).** “Effect of *Olea europaea* leaves extract on streptozotocin induced diabetes in male albino rats,” *Saudi Journal of Biological Sciences*, 26( 1); 118–128.
- Al-Brakati, A., Albarakati, A. J. A., Daabo, H. M., Baty, R. S., Salem, F. E. H., Habotta, O. A. and Amin, H. K. (2020).** Neuromodulatory effects of green coffee bean extract against brain damage in male albino rats with experimentally induced diabetes. *Metabolic brain disease*, 35, 1175-1187.
- Alexandraki, K., Piperi, C., Kalofoutis, C., Singh, J., Alaveras, A. and Kalofoutis, A. (2006).** Inflammatory process in type 2 diabetes: The role of cytokines. *Ann. NY Acad.Sci.*, 1084, 89-117
- Ali, S.J. and Shapour, H. (2013).** *Crataegus monogyna* fruit aqueous extract as a protective agent against doxorubicin-induced reproductive toxicity in male rats. *Avicenna J Phytomed.* 3(2): 159–170.

- Allain , C. C. , Poon , L. S. and Chan , C. S. (1974).** Enzymatic determination of serum total cholesterol . Clin. Chem., 20: 470-475.
- Al-Malki, A. L. and El Rabey, H. A. (2015).** The Antidiabetic Effect of Low Doses of *Moringa oleifera* Lam. Seeds on Streptozotocin-Induced Diabetes and Diabetic Nephropathy in Male Rats. *Bio. Med. Research International*, 1-13.
- Ana Angelica, H. F., Ethel, L. B., Ary, F. J. and Cristiano, M. G. (2009).** Effect of Naringin on Biochemical parameters in the streptozotocin-induced diabetic rats. *Brazilian Archives of Biology and Technology*. 52:51-59.
- AOAC (2000).** Official Methods of Analysis. 17th Edition, The Association of Official Analytical Chemists, Gaithersburg, MD, USA. Methods 925.10, 65.17, 974.24, 992.16.
- Ayala, A., Muñoz, M. F. and Argüelles, S. (2014).** “Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal,” *Oxidative Medicine and Cellular Longevity*, vol. 2014, 31 pages, .
- Barnawi, I. O., Nasr, F. A., Noman, O. M., Alqahtani, A. S., Al-zharani, M., Alotaibi, A. A. and Herqash, R. N. (2021).** Induction of apoptosis and cell cycle arrest by chloroform fraction of *Juniperus phoenicea* and chemical constituents analysis. *Open Chemistry*, 19(1), 119-127.
- Barrero, A. F., Quilez del Moral, J. F., Herrador, M. M., Akssira, M., Bennamara, A. and Akkad, S. (2004).** Oxygenated diterpenes and other constituents from Moroccan *Juniperus phoenicea* and *Juniperus thurifera* var. *Africana*. *Phytochemistry*, 65: 2507-2515.
- Bergmeyer, H.U., Horder, M. and Rey, J. (1986).** Approved recommendation on IFCC methods for the measurement of catalytic enzymes. Part 3: IFCC method for alanine aminotransferase. *J. Clin. Chem. Clin. Biochem.* 1986b; 24: 481–495.
- Beutler, E., Duron, O. and Kelly, B. M. (1963).** Improved method for the determination of blood glutathione, *The Journal of Laboratory and Clinical Medicine*, Vol. 61: 882–888.
- Bouaziz, A., Khennouf, S., Djidel, S., Baghiani, A., Dahamna, S., Amira, S. and Arrar, L. (2016).** Effects of methanolic extract of *Crataegus azarolus* L. on

plasma antioxidant capacity and biomarkers of oxidative stress in liver and kidney of rats. *International Journal of Pharmacognosy and Phytochemical Research*, 8(6), 968-976.

**Busher, J.T. (1990).** Serum albumin and globulin.clinical methods: the history, physical and laboratory examinations.3rd edition. Boston. Chapter: 101.

**Chapman, D. G., Gastilla, R. and Campbell, T. A. (1959).**Evaluation of protein in food. I. A. Method for the determination of protein efficiency ratio. *Can. j. Biochem. Physio.* 1(37): 679- 686.

**Chaudhary, S., Semwal, A., Kumar, H., Verma, H. C and Kumar, A. (2016).** *In-vivo* study for anti-hyperglycemic potential of aqueous extract of Basil seeds (*Ocimum basilicum Linn*) and its influence on biochemical parameters, serum electrolytes and haematological indices. *Biomedicine & Pharmacotherapy*, 84, 2008-2013.

**Chen, J., Zhang, X. , Huo ,D., Cao,C.,Li,Y., Liang,Y., Li. B.and Li,L. (2019).** Preliminary characterization, antioxidant and  $\alpha$ -glucosidase inhibitory activities of polysaccharides from *Mallotus furetianus*, *Carbohydrate Polymers*, . 215, pp. 307–315 .

**Chen, Z., Hu, H., Chen, M., Luo, X., Yao, W., Liang, Q., Yang, F. and Wang, X. (2020).** Association of Triglyceride to high-density lipoprotein cholesterol ratio and incident of diabetes mellitus: a secondary retrospective analysis based on a Chinese cohort study. *Lipids in Health and Disease* .19- 33.

**Cui, T., Li, J., Kayahara, H., Ma, L., Wu, L. and Nakamura, K. (2006).**Quantification of the polyphenols and triterpene acids in hawthorn by high-performance liquid chromatography. *J of Agri and Food Chem.*, 54(13):4574–4581.

**Dall,T.M., Yang, W., Halder,P., Franz,J., Byrne, E., Semilla,A.P.; Chakrabarti,R. and Stuart,B. (2016).** “Type 2 diabetes detection and management among insured adults,” *Population Health Metrics*, 14 ( 1) : 43 .

**Dar, R. , Shahnawaz, M. and Qazi, P. (2019).** General overview of medicinal plants: A review. 349-351.

**Dessoky, E.L.S., Ismail, I.A., El-Hallous, E.I. and Alsanie, W.F. (2020).**Protective Role of *Juniperus phoenicea* L. Leaves Extract against Gamma-irradiation-induced Oxidative Stress.*Pak J Biol Sci.* 23(7):922-930.

- Dewanjee, S., Chakraborty, P., Mukherjee, B., De Feo, V. (2020).** Plant-Based Antidiabetic Nanoformulations: The Emerging Paradigm for Effective Therapy. *Int. J. Mol. Sci.* , 21, 2217.
- Drupt, F. (1974):**"Depression of human serum albumin." *Farm. Boil.*, 9: 222-229.
- Elbe, H., Esrefoglu, M., Vardi, N., Taslidere, E., Ozerol, E. and Tanbek, K.(2015).** Melatonin, quercetin and resveratrol attenuates oxidative hepatocellular injury in streptozotocin-induced diabetic rats. *Hum. Exp. Toxicol.* 34: 859–868.
- Elkotby, D., Hassan, A. K. , Emad, R. and Bahgat, I.(2018).** “Histological changes in islets of Langerhans of pancreas in alloxan-induced diabetic rats following Egyptian honey bee venom treatments,” *International Journal of Pure and Applied Zoology.* 6. 1–6.
- El-Sawi, S. A. , Motawae, H. M. , El-Shabrawy, A. O. , Sleem, M. A. , Sleem, A. A. and Maamoun, M. A. N. S. (2015).** Antihyperglycemic effect of *Juniperus phoenicea* L. on alloxan-induced diabetic rats and diterpenoids isolated from the fruits,” *Journal of Coastal Life Medicine,* 3. 906–909.
- Elsawy, M. and Emara, E. (2016).**The impact of ghrelin on oxidative stress and inflammatory markers on the liver of diabetic rats. *Tanta Med J.* 44:163-169.
- Fossati, P., Prencipe, L. and Berti, G. (1980).** Use of 3, 5 dichloro-2-hydroxyl benzene sulfonic acid /4-aminophenazone chromogenic system in direct enzymatic assay of uric acid in serum and urine. *Clin. Chem.*; 26: 227-231.
- Friedwald, W. T., Leve, R. I. and Fredrickson, D. S. (1972).** Estimation of the concentration of low-density lipoprotein separated by three different methods. *Clin. Chem.*; 18: 499-502.
- Gadewar, M. M., GK, P., Mishra, P. C., Ashraf, G. M., Almashjary, M. N., Harakeh, S. and Jha, S. K. (2023).** Evaluation of Antidiabetic, Antioxidant and Anti-Hyperlipidemic Effects of *Solanum indicum* Fruit Extract in Streptozotocin-Induced Diabetic Rats. *Current Issues in Molecular Biology,* 45(2), 903-917.
- Giannopolitis, C. N. and Ries, S. K. (1977).**“Superoxide dismutase. I. occurrence in higher plants,” *Plant Physiology,* Vol. 59( 2): 309–314.



- Giribabu, N., Karim, K., Kilari, E. K. and Salleh, N. (2017).** Phyllanthus niruri leaves aqueous extract improves kidney functions, ameliorates kidney oxidative stress, inflammation, fibrosis and apoptosis and enhances kidney cell proliferation in adult male rats with diabetes mellitus. *Journal of ethnopharmacology*, 205, 123-137.
- Goyal, S. N., Reddy, N. M., Patil, K. R., Nakhate, K. T., Ojha, S., Patil, C. R. and Agrawal, Y. O. (2016).** "Challenges and issues with streptozotocin-induced diabetes—a clinically relevant animal model to understand the diabetes pathogenesis and evaluate therapeutics," *Chemico-Biological Interactions*. 244. 49-63.
- Guo, L., Jiang, B., Li, D. and Xiao, X. (2021).** Nephroprotective effect of adropinin against streptozotocin-induced diabetic nephropathy in rats: inflammatory mechanism and YAP/TAZ factor. *Drug Design, Development and Therapy*, 589-600.
- Gupta, S. C., Patchva, S., Koh, W. and Aggarwal, B. B. (2012).** Discovery of curcumin, a component of the golden spice, and its miraculous biological activities, *Clin. Exp. Pharmacol. Physiol.* 39(3): 283-99.
- Hacioglu, C., Kar, F., Kara, Y., Yucel, E., Donmez, D. B., Sentürk, H. and Kanbak, G. (2021).** Comparative effects of metformin and *Cistus laurifolius* L. extract in streptozotocin-induced diabetic rat model: oxidative, inflammatory, apoptotic, and histopathological analyzes. *Environmental Science and Pollution Research*, 28(41), 57888-57901.
- Hassan, R. M., Elsayed, M.; Kholief, T. E.; Hassanen, N. H.; Gafer, J. A. and Attia, Y. A. (2021):** Mitigating effect of single or combined administration of nanoparticles of zinc oxide, chromium oxide, and selenium on genotoxicity and metabolic insult in fructose/streptozotocin diabetic rat model. *Environmental Science and Pollution Research*, 28, 48517-48534.
- Henda, K., Kais, M., Khaled, H., Mohamed, D., Abdelfattah, E. and Nouredine, A. (2014).** In vitro anti-diabetic, anti-obesity and antioxidant properties of *Juniperus phoenicea* L. leaves from Tunisia. *Asian Pac. J. Trop. Biomed.*, 4 ( 2): S649-S655.
- Huang, Y. C. and Chen, B. H. (2023).** A Comparative Study on Improving Streptozotocin-Induced Type 2 Diabetes in Rats by Hydrosol, Extract and Nanoemulsion Prepared from Cinnamon Leaves. *Antioxidants*, 12(1), 29.

- Husdan, H. and Rapoport, A. (1968).** Estimation of creatinine by Jaffe reaction method. Clin.Chem.; 14: 222-228.
- Hussein, H. K. and Abu-Zinadah, O. A. (2010).** “Antioxidant effect of curcumin extracts in induced diabetic wister rats,” International Journal of Zoological Research, 6( 4):266–276.
- Jia ,Y., Bhuiyan, M.J., Jun, H.J., Lee,H., Hoang,M.H., Lee,H., Kim,N.,Lee.D.,Hwang,K.Y.,Hwang,B.Y.,Choi,D.W. and Lee,S.J.(2011).**Ursolic acid is a PPAR-alpha agonist that regulates hepatic lipid metabolism. Bioorg Med Chem Lett. 21:5876–80.
- Karuppusamy, A. and Thangaraj, P. (2012).** Antidiabetic activity of aqueous root extract of *Merremia tridentate* (L.) Hall in streptozotocin-induced diabetic rats, Asian Pac. J.Trop. Med. 5(3):175-179.
- Keskes, H., Mnafigui, K., Hamden, K., Damak, M., El Feki, A. and Allouche, N. (2014).** *In vitro* anti-diabetic, anti-obesity and antioxidant proprieties of *Juniperus phoenicea* L. leaves from Tunisia. Asian Pacific journal of tropical biomedicine, 4, S649-S655.
- Kota, N., Virendra Panpatil, V. R., Kaleb, B. and Polasa, V. K. (2012).** Dose-dependent the effect in the inhibition of oxidative stress and anticlastogenic potential of ginger in STZ induced diabetic rats, Food Chem., 135:2954- 2959.
- Kumar, S., Vasudeva, N. and Sharma, S. (2012).**GC-MS analysis and screening of antidiabetic,antioxidant and hypolipidemic potential of *Cinnamomum Tamala* oil in streptozotocin induced diabetes mellitus in rats, Cardiovasc.Diabetol, 11: 95-106.
- Kusunoki,J., Aragane, K, Kitamine,T., Kozono,H., Kano, K.,Fujinami, K., Kojima,K., Chiwata,T., and Sekine,Y.(2000).**“Postprandial hyperlipidemia in streptozotocin-induced diabetic rats is due to abnormal increase in intestinal acyl coenzyme A: cholesterol acyltransferase activity,” Arteriosclerosis, Thrombosis, and Vascular Biology, 20( 1); 171–178.
- Laouar, A., Klibet, F.,Bourogaa, E., Benamara, A., Boumendjel, A., Chefrou, A. and Messarah, M. (2017).** Potential antioxidant properties and hepatoprotective effects of *Juniperus phoenicea* berries against CCl4 induced

hepatic damage in rats. *Asian Pacific journal of tropical medicine*, 10(3), 263-269.

**Lo, C., Shyh-Hsiang, L., Jung-Su, C. and Yi-Wen, C. (2017).** Effects of Melatonin on Glucose Homeostasis, antioxidant ability, and adipokine Secretion in ICR Mice with Na/STZ-Induced Hyperglycemia. *Nutrients*. 9 (1187):1-13.

**Lopes-Virella, M. F., Stone, S., Ellis, S. and Collwell, J. A. (1977).** Cholesterol determination in high-density lipoprotein separated by three different methods. *Clin. Chem.*, 23(5): 882.

**Luo, D., Dong, X., Huang, J., Huang, C., Fang, G. and Huang, Y. (2021).** Pueraria lobata root polysaccharide alleviates glucose and lipid metabolic dysfunction in diabetic db/db mice. *Pharmaceutical Biology*, 59(1), 380-388.

**Luo, Y.Y., He, H.B., Zhang, M.L., Huang, X., Zhang, J., Zhou, Y., Liu, X. and Fan, N. (2014).** Serum levels of TNF- $\alpha$ , IL-6 and IL-18 elevated in chronic schizophrenic patients. *Schizophr Res*. 159:556–557.

**Mahboob, M., Rahman, M.F. and Grover, P. (2005).** Serum lipid peroxidation and antioxidant enzyme levels in male and female diabetic patients. *Singap. Med. J.* 46, 322–324.

**Makena, W., Hamman, W.O., Buraimoh, A.A., Dibal, N.I. and Obaje, S.G. (2018).** Therapeutic effects of balanitoid in streptozotocin-induced diabetic rats. *J Taibah Univ Med Sc.* 13(4):402-406.

**Mall, A. I., Maamoun, Salma, A., El-Sawi, Hemala, M., Motawae, Mohamed, A., Sleem, Abdelrahman, O., El-Shabrawy, Hawas, W., Usama, Ayman, A. and Farghaly. (2016).** Antiproliferative Effect of Extracts and Flavonoids of *Juniperus Phoenicea* L. Growing in Egypt. *Mintage journal of Pharmaceutical & Medical Sciences*, Vol. 5 (2): 1-7.

**Mandlem, V. K. K. and Annapurn, A. (2017).** “Cardioprotective role of saxagliptin through antioxidant mechanism in experimental myocardial infarction in STZ induced diabetic rats,” *Clinical & Experimental Pharmacology*, 7. 233.

**Mardiaha, B., Zakariaa, F.R., Prangdimurtia, E. and Damanik, R. (2015).** anti-inflammatory of Purple Roselle Extract in Diabetic Rats Induced by Streptozotocin. *Procedia Food Science*. 3(15 ): 182 – 189.

- Mistry, S., Dutt, K.R. and Jena,J. (2013).** Protective effect of *Sida cordata* leaf extract against CCl<sub>4</sub> induced acute liver toxicity in rats. *Asian Pac J Trop Med*, 6 (4): 280-284.
- Mohammed, A. , Koorbanally, N.A. and Islam, M.S. (2016).** Anti-diabetic effect of *Xylopiya aethiopica* (Dunal) A. Rich. (Annonaceae) fruit acetone fraction in a type 2 diabetes model of rats.*J. Ethnopharmacol.*, 180 .131-139.
- Molehin,O. R., Oloyede,O. I. and Adefegha, S. A. (2018).** “Streptozotocin-induced diabetes in rats: effects of white butterfly (*clerodendrum volubile*) leaves on blood glucose levels, lipid profile and antioxidant status,” *Toxicology Mechanisms and Methods*, 28. 1–50.
- Mostafa, D.G., Khaleel, E.F. and Abdel-Aleem, G.A.(2018).** Inhibition of the hepatic glucose output is responsible for the hypoglycemic effect of *Crataegus aronia* against type 2 diabetes mellitus in rats. *Arch Biol Sci.*70(2):277-364.
- Nacer, W., Baba Ahmed, F. Z., Merzouk, H., Benyagoub, O. and Bouanane, S. (2020).** Evaluation of the anti-inflammatory and antioxidant effects of the microalgae *Nannochloropsis gaditana* in streptozotocin-induced diabetic rats. *Journal of Diabetes & Metabolic Disorders*, 19, 1483-1490.
- Nafiu, B. A., Maung, M. C., Ni, W.,Rahela, Z. and Rahman, M. T. (2011).** Beneficial effects of ginger (*Zingiber officinale*) on carbohydrate metabolism in streptozotocin-induced diabetic rats, *British J. Nut.*, vol. 7: 1194-1201.
- Najla, O.A., Olfat, A.K., Kholoud, S.R., Enas, N.D. and Hanan, S.A. (2012).** Hypoglycemic and Biochemical Effects of *Matricaria chamomilla* Leave Extract in Streptozotocin-Induced Diabetic Rats. *Journal of Health Sciences*, 2(5): 43-48.
- Nasir, A., Abubakar, M. G., Shehu ,R. A., Aliyu, U. and Toge ,B .K. (2013).**Hepatoprotective effect of the aqueous leaf extract of *Andrographis paniculata* Nees against carbon tetrachloride – induced hepatotoxicity in rats. *Nig J Basic Appl Sci* . 21(1): 45-54.
- Navarro, J.F. and Mora, C. (2006).**Diabetes, inflammation ,proinflammatory cytokines, and diabetic nephropathy.*Sci. World J.*6, 908-917.

- Neeta, M.; Mukta, N. and Bilwa, K. (2015):** Comparative qualitative phytochemical analysis of *Sesamum indicum* L. 2 172-181.
- Nwagha, U.I.(2010).** Atherogenic index of plasma as useful predictor of cardiovascular risk among postmenopausal women in Enugu, Nigeria. *African Health Sciences*. 10(3): 248-252.
- Okechukwu, P.N., Ndyebura, A.W., Chiang, C.N and Akowuah, G.A. (2013).** Effect of Standardized Extract of *Cosinium fenestratum* Stem Bark on Liver and Kidney Function Parameters in Streptozotocin-Induced Diabetic Rats. *Journal of Acute Disease*, 2(3): 201-206.
- Okoduwa, S. I., Umar, I. A., James, D. B. and Inuwa, H. M. (2017).** Anti-diabetic potential of *Ocimum gratissimum* leaf fractions in fortified diet-fed streptozotocin treated rat model of type-2 diabetes. *Medicines*, 4(4), 73.
- Ono, Y., Hattori, E., Fukaya, Y., Imai, S. and Ohizumi, Y. (2006 ).** Anti-obesity effect of *Nelumbo nucifera* leaves extract in mice and rats. *J. Ethnopharmacol*, 106: 238-244.
- Othman, G. Q. and Mustafa, T. A. (2017).** The liver protective role of hawthorn (*Crataegus* sp.) in hypertriglyceridemic induced rats. *Polytechnic Journal*, 7(2), 111-118.
- Özcan, M.A., Haciseferoğulları, H., Marakoğlu, T. and Arslan, D. (2005).** Hawthorn (*Crataegus* spp.) fruit: Some physical and chemical properties. *Journal of Food Engineering* 69(4):409-413.
- Patton, C. J. and Crouch S. R. (1977).** Enzymatic colorimetric method for determination of urea in serum. *Anal. Chem.*, 49: 464-465.
- Pirmoghani, A., Salehia, I., Moradkhani, S., Karimia, S.A. and Salehi, S. (2019).** Effect of *Crataegus* extract supplementation on diabetes induced memory deficits and serum biochemical parameters in male rats. *J. IBRO Reports* . 7, 90-96.
- Qin, C., Xia, T., Li, G., Zou, Y.; Cheng, Z. and Wang, Q. (2019).** Hawthorne leaf flavonoids prevent oxidative stress injury of renal tissues in rats with diabetic kidney disease by regulating the p38 MAPK signaling pathway. *Int J Clin Exp Pathol* .12(9):3440-3446.

- Rababa'h, A. M., Al Yacoub, O. N., El-Elimat, T., Altarabsheh, S., Deo, S., Al-Azayzih, A. and Alzoubi, K. H. (2020).** The effect of hawthorn flower and leaf extract (*Crataegus Spp.*) on cardiac hemostasis and oxidative parameters in Sprague Dawley rats. *Heliyon*, 6(8), e04617.
- Raina, R., Verma, P. K., Peshin, R. and Kour, H. (2019).** Potential of *Juniperus communis L* as a nutraceutical in human and veterinary medicine. *Heliyon*, 5(8), e02376.
- Ramesh, B.K., Maddirala, D.R., Vinay, K.K., Shaik, S.F., Tiruvenkata, K.E.G., Swapna, S., Ramesh, B., Rao, C.A.(2010).** Antihyperglycemic and antihyperlipidemic activities of methanol:water (4:1) fraction isolated from aqueous extract of *Syzygium alternifolium* seeds in streptozotocin induced diabetic rats. *Food. Chem. Toxicol.* 48, 1078–1084
- Reeves, P. G., Nielson, F. H. and Fahmy, G. C. (1993).** Reports of the America Institute of Nutrition, adhoc wiling committee on reformulation of the AIN93. Rodent Diet. *J.Nutri.*, 123, 1939-1951.
- Roy, S. E. (1970):**"Colorimetric determination of serum alkaline phosphatase". *J. Clin. Chem.*, 16:431-432.
- Sadri, H., Goodarzi, M., Salemi, Z.and Morteza, S.(2017).**antioxidant Effects of Biochanin a inStreptozotocin Induced Diabetic Rats. *Braz.arch. biol. Technol.* 60:11-19.
- Salma, A.E.M., Eman, R.Z., Heba, M.I.A. and Mahmoud, S.A.(2017).** Therapeutic effects of aqueous, methanol and ethanol extracts of Egyptian *Artemisia herba-alba* in STZ-induced diabetic neuropathy in rats. *J Appl Pharm Sci.* 7:180–187.
- Saswata, B., Hariom, S. and Tapan, K. C. (2013).** Evaluation of Anti-Diabetic and Anti-Hyperlipidemic Potential of Methanolic Extract of *Juniperus communis (L.)* in Streptozotocinnicotinamide Induced Diabetic Rats. *Int. J. Pharm. Bio. Sci.*, 4(3): 10 – 17.
- Sekkin, S., Ipek, E., Boyacioğlu, M. and Kum ,C.(2015).**DNA protective and antioxidative effects of melatonin in streptozotocin-induced diabetic rats. *Turkish Journal of Biology.* 39: 932-940.

- Senthilkumar, G.P., Arulselvan, P., Sathishkumar, D. and Subramanian, S.P.(2006).** Antidiabetic activity of fruits of *Terminalia chebula* on streptozotocin induced diabetic rats. *J. Health Sci.* 52, 283–291.
- Shalaby, M.A. and Hamowieh, A.R. (2010).** Safety and efficacy of *Zingiber officinale* roots on fertility of male diabetic rats. *Food Chem. Toxicol.*; 48:2920-2924.
- Siest, G., Henny, F. and Schiele F. (1981).** enzymatic determination of glucose. *Interpret.Exam. Lab.* 2:206-213.
- Snedecor, G. W. and Cochran, W. G. (1989).** *Statistical Methods.* 8th ed., Iowa State University Press, Ames, Iowa 50014, USA.
- Soliman, G.Z.A. (2013).** Effect of Vitamin C and/or Vitamin E on Kidney, Liver and Brain Functions of Streptozotocin-Induced Diabetic Rats. *The Egyptian Journal of Hospital Medicine*, 53: 799–808.
- Sonnenwirth, A. and Jaret, L. (1980).** *Grad Wholes Clinical Laboratory Methods and Diagnosis.* 18 ed Mosby, London: 258-259.
- Suanarunsawat, T. ; Anantasomboon, G. and Piewbang, C. (2016):** Anti-diabetic and anti-oxidative activity of fixed oil extracted from *Ocimum sanctum L.* leaves in diabetic rats *Exp. Ther. Med.* 11. 832-840.
- Suryanarayana, N. P., Saraswat, M., Mrudula, T., Krishna, T. P., Krishnaswamy, K. and Bhanuprakash, G. R. ( 2005).** Curcumin and turmeric delay streptozotocin-induced diabetic cataract in rats, *Invest. Ophthalmol. Vis. Sci.*, 46: 2092- 2099.
- Tebboub, I. and Kechrid, Z. (2021).** Effect of curcuma on zinc, lipid profile and antioxidants levels in blood and tissue of streptozotocin-induced diabetic rats fed zinc deficiency diet. *Archives of Physiology and Biochemistry.* 127 162-169.
- Triki, R., Boughediri, K., Chelaghmia, M. L. and Kechrid, Z. (2022).** Effect of *Crataegus Azarolus* on blood glucose, lipid profile and antioxidant status in streptozotocin diabetic rats fed zinc deficient diet.1-27.
- Triki, R., Boughediri, K., Chelaghmia, M. L. and Kechrid, Z. (2022).** Effect of *Crataegus Azarolus* on blood glucose, lipid profile and antioxidant status in streptozotocin diabetic rats fed zinc deficient diet.1-27.



- Trinder, P. and Ann, S. (1969).** Enzymatic Colorimetric test with lipid clearing factor to determine triglycerides. *Clin .Biochem.*, 6: 24-27.
- Uotila, M., Ruosiahti, E. and Engrall, E. (1981).** Enzyme Linked Immunoassays. *Journal of Immunological Method*, 42, 11-15.
- Wahba, N. S., Shaban, S. F., Kattaia, A. A. and Kandeel, S. A. (2016).** "Efficacy of zinc oxide nanoparticles in attenuating pancreatic damage in a rat model of streptozotocin-induced diabetes," *Ultrastructural Pathology*, . 40( 6); 358–373.
- Wang, S. Z., Wu, M., Chen, K. J., Liu, Y., Sun, J., Sun, Z. and Liu, L. T. (2019).** Hawthorn extract alleviates atherosclerosis through regulating inflammation and apoptosis related factors: an experimental study. *Chinese journal of integrative medicine*, 25, 108-115.
- Xu ,Y.J., Tappia, P.S., Neki, N.S. and Dhalla, N.S.(2014a).**Prevention of diabetes induced cardiovascular complications upon treatment with antioxidants. *Heart Fail Rev.* 19: 113–121.
- Xu, Z., Wei, Y., Gong, J., Cho, H., Park, J. K., Sung, E. R. and Duh, E. J. (2014b).** NRF2 plays a protective role in diabetic retinopathy in mice. *Diabetologia*, 57, 204-213.
- Yahyaoui, A., Arfaoui, M. O., Rigane, G., Hkir, A., Amari, K., Ben Salem, R. and Ammari, Y. (2019).** Investigation on the chemical composition and antioxidant capacity of extracts from *Crataegus azarolus L.*: effect of growing location of an important Tunisian medicinal plant. *Chemistry Africa*, 2, 361-365.
- Yallow, R. and Bauman, W. A. (1983).** Plasma insulin in health and disease. In *Diabetes Mellitus: Theory and Practice*. Vol. 15. New York: Excerpta Medica. p. 119-20.
- Yang, D. and Hyung-Sub, K(2018).** anti-Diabetic Effect of Cotreatment with Quercetin and Resveratrol in Streptozotocin-Induced Diabetic Rats. *Biomol Ther (Seoul)*. 26(2): 130–138.
- Yang, L., Wang, C.Z., Ye, J.Z. and Li, H.T.(2011).** Hepatoprotective effects of polyphenols from Ginkgo biloba L. leaves on CCl<sub>4</sub>-induced hepatotoxicity in rats *Fitoterapia*, 82 (6) 834-840.

- Yoshioka, T., Kawada, K., Shimada, T. and Mori, M. (1979).**“Lipid peroxidation in maternal and cord blood and protective mechanism against activated-oxygen toxicity in the blood,” *The American Journal of Obstetrics and Gynecology*. 135( 3): 372–376.
- Zafar, M. and Naqvi,S. N.(2010).** “Effects of STZ-induced diabetes on the relative weights of kidney, liver and pancreas in albino rats: a comparative study,” *International Journal of Morphology*. 28; 135–142.
- Zafar, M., Naqvi, S.N., Ahmed, M. and Kaimkhani, Z.A.(2009).** Altered Liver Morphology and Enzymes in Streptozotocin Induced Diabetic Rats.*International Journal of Morphology*, 27(3): 719-725.
- Zhang, B., Yue, R., Chen, Y.,Yang, M., Huang, X., Shui, J., Peng, Y. and Chin, J. (2019)** .Gut Microbiota, a Potential New Target for Chinese Herbal Medicines in Treating Diabetes Mellitus. *Evidence-BasedComplementary and Alternative Medicine* 2019 2634898.
- Zhang, C. Y. and Tan, B. K. (2002).**Antihyperglycemic and anti-oxidant properties of *Andrographis paniculate* in normal and diabetic rats, *Clin. Exp.* 27: 358-363
- Zhang, J., Chai, X., Zhao, F., Hou, G. and Meng, Q. (2022).** Food Applications and Potential Health Benefits of Hawthorn. *Foods*, 11(18), 2861.
- Zhang, Y., Feng, F., Chen, T., Li, Z. and Shen, Q.W. (2016).** Antidiabetic and antihyperlipidemic activities of *Forsythia suspensa* (Thunb.) Vahl (fruit) in streptozotocin-induced diabetes mice *J. Ethnopharmacol.*, 192 . 256-263.
- Zhao,J.,Liu,T.,Xu,F.,You,S.,Xu,F.,Li,C. and Gu,Z.(2016).**Anti-arthritis effects of total flavonoids from *Juniperus sabina* on complete freund's adjuvant induced arthritis in rats.*Phcog Mag* .12( 47):178-183.

## تأثير المستخلصات الكحولية لنبات العرعر الفينيقي و الزعرور الشائع على مرض السكر المحدث بالستربتوزوتوسين في فئران التجارب

نيفين مصطفى زعيمة

قسم التغذية وعلوم الأطعمة - كلية الاقتصاد المنزلي - جامعة الأزهر - مصر

### المستخلص العربي

مرض السكري عبارة عن مجموعة غير متجانسة من الاضطرابات التي تتميز بنقص إفراز الأنسولين أو وجود حساسية للأنسولين. لذلك، صممت الدراسة الحالية لمعرفة التأثير المضاد للمستخلصات الكحولية لأوراق العرعر الفينيقي والزعرور الشائع علي ارتفاع سكر الدم في الفئران المصابة بمرض السكري التي يسببها الستربتوزوتوسين. تم استخدام 30 فأر من سلالة الألبينو تتراوح أوزانهم بين  $(200 \pm 10 \text{ جم})$ . وتم تقسيم الفئران إلى 6 مجموعات متساوية مجموعة (1) تركت أحد المجموعات كمجموعة ضابطة سالبه، المجموعة (2) تم تغذيتها علي الغذاء القياسي كمجموعة ضابطة موجبة، (3 و 4) تمت معالجتهم بمستخلص العرعر الفينيقي (200 و 400 مجم لكل كجم من وزن الجسم) على التوالي، (5 و 6) تمت معالجتهم بمستخلص الزعرور الشائع (200 و 400 مجم لكل كجم من وزن الجسم) على التوالي لمدة أربعة أسابيع. في نهاية فترة التجربة تم حساب التقييم البيولوجي. وتم تقدير الجلوكوز والأنسولين، وظائف الكبد، إنزيمات القلب، ووظائف الكلى، دهون الدم وعامل النخر في السيرم. وتم تقدير بعض العوامل المؤكسدة ومضادات الأكسدة في أنسجة البنكرياس. كما تم تحليل التركيب الكيميائي للأعشاب. أظهرت النتائج أن المستخلصات الكحولية لأوراق العرعر الفينيقي و الزعرور الشائع أدت إلي زيادة في التقييم البيولوجي وتحسين في وظائف الكبد، ووظائف الكلى، وتحسن مستوى الدهون، وانخفاض نسبة الجلوكوز في الدم، وإنزيم القلب، و MDA ولكن زيادة في مستويات الأنسولين و GSH و SOD مقارنة بالفئران المصابة بمرض السكري وغير المعالجة. لذلك أوصت الدراسة بأن تناول مستخلص أوراق العرعر الفينيقي والزعرور الشائع يمكن أن تقلل من مستوي سكر الدم والمضاعفات الناتجة عن ارتفاع سكر الدم.

**الكلمات المفتاحية:** العرعر الفينيقي، الزعرور الشائع، الستربتوزوتوسين، سكر الدم، الأنزيمات المضادة

للأكسدة.